


Amendments to the Claims:


Please amend the claims as follows (the changes in these claims are shown with ~~strikethrough~~ for deleted text and underlines for added text). A complete listing of the claims is listed below with proper claim identifiers.

Listing of Claims:

1. (Currently Amended) A composition, comprising:
a quaternary ammonium compound of formula (I)



a phosphate ester of formula (II);



wherein R^1 , R^2 , R^3 , R^4 are independently selected from the group consisting of alkyl, alkenyl and alkynyl groups;


X^- is selected from the group consisting of halide and sulfate; and

R^5 is a polyoxyalkylated branched alcohol, and R^6 , and R^7 are independently selected from the group consisting of hydrogen, a hydrocarbyl group, and a polyoxyalkylated alcohol.

2. (Original) The composition of claim 1, wherein R^1 and R^2 contain from 1 to 6 carbon atoms; and R^3 and R^4 contain from 7 to 20 carbon atoms.

3. (Original) The composition of claim 1, wherein R^1 and R^2 contain from 1 to 5 carbon atoms; and R^3 and R^4 contain from 7 to 15 carbon atoms.

4. (Original) The composition of claim 1, wherein R^1 and R^2 contain from 1 to 3 carbon atoms; and R^3 and R^4 contain from 8 to 12 carbon atoms.
5. (Original) The composition of claim 1, wherein R^1 and R^2 are decyl; and R^3 and R^4 are methyl.
6. (Previously presented) The composition of claim 5, wherein X^- is a halide.
7. (Previously presented) The composition of claim 5, wherein X^- is chloride.
8. (Currently Amended) The composition of claim 1, wherein R^5 is a ~~polyoxyalkylated alcohol~~ of comprises from 2 to 500 carbon atoms.
9. (Currently Amended) The composition of claim 8, wherein R^5 the ~~polyoxyalkylated alcohol~~ comprises an alcohol portion of from 1 to 20 carbon atoms.
10. (Currently Amended) The composition of claim 8, wherein R^5 the ~~polyoxyalkylated alcohol~~ comprises an alcohol portion of from 6 to 14 carbon atoms.
11. (Original) The composition of claim 8, wherein R^6 and R^7 are hydrogen.
12. (Original) The composition of claim 1, wherein the phosphate ester is poly(oxy-1,2-ethandiyl) tridecyl hydroxy phosphate.
13. (Previously presented) The composition of claim 1, further comprising a thiocarbonyl compound of formula (III)



(III);

wherein R^8 is selected from the group consisting of metal ion, ammonium ion, hydrocarbyl, and heterohydrocarbyl;


X and Y are independently selected from the group consisting of oxygen and sulfur such that at least one of X and Y is sulfur;

Z is selected from the group consisting of OR^9 and $NR^{10}R^{11}$; and


R^9 , R^{10} , and R^{11} are independently selected from the group consisting of hydrocarbyl and heterohydrocarbyl.

14. (Original) The composition of claim 13, wherein X is sulfur.
15. (Original) The composition of claim 14, wherein Z is $NR^{10}R^{11}$.
16. (Original) The composition of claim 15, wherein R^{10} and R^{11} are independently hydrocarbyl groups of from 1 to 10 carbon atoms.
17. (Original) The composition of claim 15, wherein R^{10} and R^{11} are independently hydrocarbyl groups of from 1 to 5 carbon atoms.
18. (Original) The composition of claim 16, wherein Y is sulfur.
19. (Original) The composition of claim 18, wherein R^8 is a metal ion.
20. (Original) The composition of claim 13, wherein the thiocarbonyl compound is potassium dimethyl dithiocarbamate.
21. (Original) The composition of claim 1, further comprising a solvent.
22. (Original) The composition of claim 1, further comprising at least one additive selected from the group consisting of a supplemental corrosion inhibitor, a scale inhibitor, a surfactant, a biocide, a foamer, and an oxygen scavenger.

23. (Previously presented) A composition, comprising:
a quaternary ammonium compound of formula (I)




wherein R^1 , R^2 , R^3 , R^4 are independently a hydrocarbyl group;
a phosphate ester of formula (II);



wherein X^- is selected from the group consisting of halide and sulfate; and

R^5 , R^6 , and R^7 are independently selected from the group consisting of hydrogen, a hydrocarbyl group, and a polyoxyalkylated alcohol; and
a thiocarbonyl compound of formula (III);



wherein R^8 is selected from the group consisting of metal ion, ammonium ion, hydrocarbyl, and heterohydrocarbyl;

X and Y are selected from the group consisting of oxygen and sulfur, such that at least one of X and Y is sulfur; and


R^{10} and R^{11} are independently selected from the group consisting of hydrocarbyl and heterohydrocarbyl.

24. (Original) The composition of claim 23, wherein
R¹ and R² are independently a hydrocarbyl group of from 1 to 6 carbon atoms;
R³ and R⁴ are independently a hydrocarbyl group of from 7 to 20 carbon atoms;
R⁵ is a polyoxyalkylated alcohol of from 2 to 500 carbon atoms;
R⁶ and R⁷ are independently hydrogen or a hydrocarbyl group of from 1 to 20 carbon atoms;
X is sulfur; and
R¹⁰ and R¹¹ are independently hydrocarbyl groups of from 1 to 10 carbon atoms.
25. (Original) The composition of claim 23, wherein the quaternary ammonium compound is didecyl dimethyl ammonium chloride; the phosphate ester is poly(oxy-1,2-ethandiyl) tridecyl hydroxy phosphate; and the thiocarbonyl compound is potassium dimethyl dithiocarbamate.
26. (Original) The composition of claim 23, further comprising a solvent.
27. (Original) The composition of claim 26, further comprising at least one additive selected from the group consisting of a supplemental corrosion inhibitor, a scale inhibitor, a surfactant, a biocide, a foamer, and an oxygen scavenger.
28. (Currently Amended) The composition of claim 27, wherein
the quaternary ammonium compound is present at ~~1-95~~ 92% by weight;
the phosphate ester is present at ~~1-95~~ 92% by weight;
the thiocarbonyl compound is present at ~~1-95~~ 92% by weight;
the solvent is present at 5-95% by weight; and
the at least one additive is present at ~~1-95~~ 92% by weight.


29. (Previously presented) The composition of claim 27, wherein
the quaternary ammonium compound is present at 1-50% by weight;
the phosphate ester is present at 1-50% by weight;
the thiocarbonyl compound is present at 1-50% by weight;
the solvent is present at 20-80% by weight; and
the at least one additive is present at 1-50% by weight.
30. (Previously presented) The composition of claim 27, wherein
the quaternary ammonium compound is present at 1-20% by weight;
the phosphate ester is present at 1-20% by weight;
the thiocarbonyl compound is present at 1-20% by weight;
the solvent is present at 50-75% by weight; and
the at least one additive is present at 1-20% by weight.
31. (Original) The composition of claim 27, wherein the quaternary
ammonium compound, the phosphate ester, and the thiocarbonyl compound are
present at a 1:1:1 ratio by volume.
32. (Original) A method of inhibiting corrosion of iron and ferrous base
materials, comprising:
contacting a material with the composition of claim 1.
33. (Original) A method of inhibiting corrosion of iron and ferrous base
materials, comprising:
contacting a material with the composition of claim 23.
34. (Original) A method of inhibiting corrosion of iron and ferrous base
materials, comprising:
contacting a material with the composition of claim 25.

35. (Currently Amended) A method of making a corrosion inhibitor, comprising

combining a quaternary ammonium compound of formula (I)



with a phosphate ester of formula (II)




wherein R^1 , R^2 , R^3 , R^4 are independently selected from the group consisting of alkyl, alkenyl and alkynyl groups;

X^- is selected from the group consisting of halide and sulfate; and


R^5 is a polyoxyalkylated branched alcohol, and R^6 , and R^7 are independently selected from the group consisting of hydrogen, a hydrocarbyl group, and a polyoxyalkylated alcohol.

36. (Previously presented) A method of making a corrosion inhibitor, comprising


combining a quaternary ammonium compound of formula (I)



with a phosphate ester of formula (II)



and further with a thiocarbonyl compound of formula (III)



wherein R^1 , R^2 , R^3 , R^4 are independently a hydrocarbyl group;

X^* is selected from the group consisting of halide and sulfate;

R^5 , R^6 , and R^7 are independently selected from the group consisting of hydrogen, a hydrocarbyl group, and a polyoxyalkylated alcohol;

R^8 is selected from the group consisting of metal ion, ammonium ion, hydrocarbyl, and heterohydrocarbyl;

X and Y are independently selected from the group consisting of oxygen and sulfur such that at least one of X and Y is sulfur;

Z is selected from the group consisting of OR^9 and $\text{NR}^{10}\text{R}^{11}$; and

R^9 , R^{10} , and R^{11} are independently selected from the group consisting of hydrocarbyl and heterohydrocarbyl.